

Instructions for Use

GKE Biological Indicators

1. Spore Strips and Spore Discs

ArtNo.*	Product Code	Quantity	Pop.	Sterilization	Content Carrier	Biological indicator	Incubation
221-601		100		process		indicator	temperature
221-605	B-E-H-SS-10-6	500	-		Spore strips filter	B. atrophaeus	33-37°C
221-610	D E 11 00 10 0	1.000	10 ⁶	Ethylene oxide,	paper		
221-611		110	- 10	Dry heat	Spore discs filter		
221-612	B-E-H-SD-10-6	100			paper		
223-501		100					55-60°C
223-505	B-S-SS-10-5	500	10 ⁵	Steam			
223-510		1.000			Spore strips filter paper		
223-601		100		Steam, Formaldehyde**			
223-605	B-S-F-SS-10-6	500	10 ⁶				
223-610		1.000		Formaldenyde			
332-407	B-V-ST-SS-10-4	100			Spore strips stainless	-	
332-415	B-V-ST-DIS-SP-10-4	110	10 ⁴		Spore discs stainless		
332-417	B-V-ST-DIS-SP-10-4	100			steel		
332-507	B-V-ST-ST-10-5	100	105		Spore strips stainless		
332-515	B-V-ST-DIS-SP-10-5	110			Spore discs stainless	Geob. Stearo- thermophilus	
332-517	B-V-ST-DIS-SP-10-5	100			steel		
332-601	B-V-P-SS-10-6	100			Spore strips PET		
332-604	2 1 1 00 10 0	40	-				
332-602	B-V-G-SS-10-6	100	-		Spore strips glass		
332-605		40	-	Hydrogen	fiber		
332-603	B-V-T-SS-10-6	100	-	peroxide /	Spore strips Tyvek°		
332-606		40	-	Plasma			
332-607	B-V-ST-SS-10-6	100 40	-		Spore strips stainless steel		
332-608 332-612		100	10 ⁶		31661		
332-612	B-V-P-DIS-SP-10-6	110	-		Spore discs PET		
332-614		100	-		Spore discs glass		
332-611	B-V-G-DIS-SP-10-6	110	-		fiber		
332-618		100	-				
332-613	B-V-T-DIS-SP-10-6	110	-		Spore discs Tyvek®		
332-617		100	-		Spore discs stainless		
332-615	B-V-ST-DIS-SP-10-6	110	-		steel		

2. Growth medium, diameter: 16.1 mm (accessories to incubate biological indicators)

Art No.*	Product Code	Quantity/ pack	Growth media	Color change of media after sterilization and incubation		Biological	Sterilization
			additive	Sterile	Non- sterile	indicator	process tested
221-010	B-E-H-CM	10	pH-indicator	Green	Yellow-	B. atrophaeus	Ethylene oxide, Dry heat
221-100		100			orange		
223-010	B-S-V-CM	10	pi i-indicatoi	Purple	Yellow- green	G. stearothermophilus	Steam, Hydrogen peroxide
223-100	B-3-V-CM	100					

^(*) To all article numbers a 3-digit alpha code is added. The additional letter code refers to the language and/or customized version. It is only added on the outside label; the inside of

Application

The GKE biological indicators (spore strips and discs) are used for validation and monitoring of sterilization and decontamination processes. For evaluation, a microbiological laboratory is required. GKE offers specific growth media with a pH-indicator. The growth medium has been developed to incubate G. stearothermophilus and B. atrophaeus biological indicators to easily monitor growth. During the growth of the bacteria the pH-value of the medium is changing the colour allowing a quicker evaluation of biological indicators.

Product Description

Spore Strips:

The GKE biological spore strips consist of a contaminated carrier of 6 x 38 mm in a glassine packaging that protects the spore strip from re-contamination after sterilization. For steam, ethylene oxide and formaldehyde filter paper is used as a carrier while hydrogen peroxide $(H_2O_2/plasma)$ strips uses different carrier materials. The spore strips for hydrogen peroxide sterilization processes are individually packaged or

the pack is identical to the article numbers on the above table.

(**) The D-value for steam is specified in the certificate, not the D-value for formaldehyde (LTSF). The D-value for LTSF can be determined at extra cost (Art.no. 223-998).

alternatively packaged together. The glassine packaging has to be opened in a microbiological laboratory where the indicator strip is aseptically transferred into the GKE Steri-Record® growth medium.

Spore Discs:

The spore discs (diameter 7 mm) consist of *B. atrophaeus* or *G. stearothermophilus* spores that are immobilized on different carriers. The spore discs are packaged together or individually in a blister box.

Growth Medium:

The European Pharmacopeia (EP) requires an incubation time of

7 days since alive but damaged spores need a longer time for growth. The GKE growth medium is optimized so that an incubation time of 7 days is not necessary. The reduced incubation times (RIT) are measured according to EN ISO 11138-8. The growth speed heavily depends on the growth medium used. The incubation times mentioned in our directions for use can only be assured if the GKE growth medium is used.

The glass tube with the length of 7,6 cm and an inner diameter of 16,1 mm, contains 8 ml sterile CASO bouillon optimized for a quick G. stearothermophilus and B. atrophaeus bacteria growth. The medium contains different pH-indicators for a quicker readout. The medium volume has been optimized for spore strips of 6 x 38 mm.

Performance Characteristics

All GKE biological indicators are produced in accordance with the EN ISO 11138-1 standard and the current European and United States Pharmacopoeia (EP+USP). Population and Dvalue are measured and certified for each batch documented in the enclosed certificate.

For hydrogen peroxide a standard does not yet exist since no appropriate resistometers/test sterilizers are available to measure D-values. Therefore, most manufacturers have their own test procedures that are not comparable. GKE has developed a test method for hydrogen peroxide in liquid $\rm H_2O_2$ solution. This test method has been presented to the Standard committee and will be published later and has the advantage that resistances can be compared throughout the world.

For information on carrying out the population determination, please refer to GKE Technical Information TI 730-067 EN.

Handling Information

1. Use of

Spore Strips:

A minimum of two GKE spore strips indicators shall be used to monitor each sterilizer cycle. Place them in the most difficult location in the sterilizer chamber. Alternatively, the spore strips can also be used in process challenge devices (PCD). For this specific use the spore strips have to be taken out of the package and put into the PCD. Place the PCD in the lower section of the sterilizer close to a door. After sterilization, the strips must not be removed and the closed PCD shall be opened aseptically in a microbiological lab.

Spore Discs:

To decontaminate isolators and rooms/chambers hydrogen peroxide processes can be used. For monitoring spore discs are positioned at the most critical areas e.g., on apparatuses/devices.

- a) After sterilization or decontamination, aseptically transfer the spore strip or disc into sterile tryptic broth. Preferentially use GKE growth medium, containing a pHindicator for fast and easy evaluation.
 - b) After LTSF sterilization processes, prior to incubation, inactivate remaining FO residues on the strips according to EN ISO 11138-5.
- 3. It is recommended to run a vitality test. Use an unsterilized control strip to guarantee the vitality of the strips and to check quality of the media. For the negative control incubate the culture medium without spore strip to exclude contamination of the media or tube. Growth of the vitality test should be evident in 2 days latest. The tube without spore strip shall not change color.
- Incubate sterilized spore strips and discs for a maximum of 7 days at 55-60°C for G. stearothermophilus and 33-37°C for B. atrophaeus and B. subtilis.
- 5. Observe all tubes for growth daily.
- Record results.
- If a positive result is observed, repeat the test procedures, placing several indicators throughout the sterilizer. If a positive result is observed again, the sterilizer has to be examined for malfunction.
- After a repair, repeat the test with several indicators and proceed only to use the sterilizer if all tests show a negative result.

Safety Precautions

Standard growth medium for steam sterilization processes should only be used in accordance with GKE Technical Information TI 730-041 when evaluating biological indicators for testing NTDF (low temperature steam formaldehyde) as this could lead to false negative results (see EN ISO 11138-5).

Storage and Disposal

- Store all spore strips in the original package between 5-30°C with a humidity of 5-80% RH away from sterilizing agents. Store the growth medium in a dark place between 5-30°C.
- 2. Biological indicators should not be used after expiry date.
- 3. Dispose biological indicators with normal waste. Sterilize biological indicators before disposal.

For further technical details please contact your local dealer or the GKE application laboratory. We will assist you with any technical questions. Also visit our website www.gke-healthcare.com for more information.

760-086 EN V11 07/2024